

Diversity and Status of Butterflies in Maitum Village, Tandag, Surigao del Sur, Philippines

RAY KRISTOFFER CASIO RAMIREZ

crinoidz@yahoo.com/ray.ramirez.biologist@gmail.com

ALMA B. MOHAGAN

almohagan@gmail.com

Central Mindanao University
Philippines

Date Submitted: June 29, 2011

Final Revision Accepted: July 10, 2011

Abstract - Diversity and assessment of butterflies of Maitum, Tandag, Surigao del Sur were conducted to provide information on the species composition, diversity and status of butterflies in dipterocarp forest and agro ecosystems. A total of 104 species of butterflies belonging to 68 genera and 5 families were documented. Species richness of the butterflies was higher in the dipterocarp forest (89) than the agro ecosystem (51). The dipterocarp forest had greater mean individual (4.173) than the agro ecosystem (3.712). The diversity level was fair in dipterocarp forest ($H' = 1.638$) as compared to the agroecosystem area with low level of diversity ($H' = 1.369$). Dendrogram for similarity of species composition showed a fairly low species between the two study stations, this suggests the importance of the two habitats for the conservation of butterflies in Maitum Village, Tandag, Surigao del Sur. Assessment of status showed that 57 species (54.8 %), were common 16 (15.3%) were rare, 12 (11.5%) were rare Philippine endemics, 6 (5.7%)

were common Philippine endemics, 2 (1.9%) common Mindanao endemics, 1 (.96%) rare Mindanao endemic, 1 (.96%) very rare eastern Mindanao endemic, and 9 (8.6%) were undetermined. These results suggest that Maitum, Tandag, Surigao del Sur is the home of 104 species of butterflies and the dipterocarp forest is the home of diverse and endemic species of butterflies for conservation.

Keywords - Diversity, Status, Conservation, Butterflies, Maitum Village.

INTRODUCTION

Tandag City is in the north-eastern coast of Mindanao. The city is the provincial capital of Surigao del Sur and one of the newly proclaimed cities in the country (United Tourist Promotions, 2008). It lies on the eastern part of Surigao del Sur which is part of Region XIII or Caraga Region. It has a total land area of 29, 173 hectares. Of these 5, 911 has. are alienable and disposable and 25, 709 are forestlands. Tandag falls under the Type II climate in the country, which is characterized by rainfall distributed throughout the year, with a negligible short dry season (MPDO Tandag Executive Summary 2004).

A study on mammal diversity by Wargues et al. (2007), taken from the WCSP 18th Annual Biodiversity Symposium "Biodiversity in the Philippine Uplands" Symposium Manual, reported a total of 28 species in Tandag, Surigao del Sur. No study on the diversity of butterflies has been reported on the area yet. This study focused on the diversity of butterflies in Maitum, Tandag, Surigao del Sur.

MATERIALS AND METHODS

Description of Study Site

This study was conducted in selected locations in Maitum, Tandag, Surigao del Sur (Figs. 1 & 2). The site is partially inhabited and is composed of two vegetation types: Agroecosystem and Dipterocarp Forests. The topography is generally plain, rolling and gently sloping. From Tandag City proper, the study site can be reached by

motorcycles in approximately 30 minutes. The study was done from April to December 2010. Maitum Village has a total land area of 1,327.37 hectares. Tandag falls under the Type II climate in the country, which is characterized by rainfall distributed throughout the year, with a negligible short dry season (MPDO Tandag Executive Summary 2004).

Establishment of Study Stations

Two study stations were established. The following were:

Station 1: Agroecosystem area (Fig.3)

Station 2: Dipterocarp forest area (Fig.4)

Sampling of butterflies was done in the above established stations of the study site.

Sampling Techniques

The following sampling techniques were employed:

1. Transect walk sampling

This was done by walking three hours from 7:00AM to 10:00AM through the 2 km transects in each study station (Figs. 5 & 6), adapted from Mohagan, and Treadaway 2010. All butterflies seen along the 2 km transect were collected, counted and listed. Butterflies seen 5 m ahead were included in the collection (Pollard and Yates 1995).

A total of 6 weeks were devoted for sampling butterflies last summer 2010 at 3 weeks per station. Another 2 weeks was allotted for sampling in December 2010 at 1 week sampling time per station. Collections of voucher specimens were limited to 3 to 5 individuals only per species. Sample duplicates were released after each transect walk.

2. Opportunistic Sampling

To enhance species richness, Opportunistic Sampling was employed. This method was used to document butterflies outside the transect. Areas around the transect were scanned for butterfly species to enrich listings (Chiong 2007). A total of 6 weeks were devoted for

sampling butterflies last summer at 3 weeks per station. Another 2 weeks was allotted for sampling in December at 1 week sampling time per station.

Collection of Specimens

Collection of butterflies was done using insect catching nets made of fine silk material to avoid wing damage. This was enhanced also by employing baits like fruit bait made of over ripe banana or any available fruit (Fig. 7). Brightly colored clothes were also used during sampling to attract butterflies to the researcher's location (Fig. 8). Butterflies collected were deposited in CMU Museum.

Classification and Identification

Classification of butterflies was done using references like books, journals photographs of previously identified specimens. Examples of these include: Butterflies of the World by Schroeder and Treadaway, Butterflies of the South East Asian Islands, The Zoological Society of London Journal, and Field Identification Guides by other authors.

Analysis of Data (Biodiversity Assessment)

Using BioPro software version 2.0, the following biodiversity indices was determined: Shannon-Weiner Diversity Index, Species Richness, Species Abundance and Bray-Curtis Similarity Index of Species Composition.

Assessment of Status

The national status and distribution of collected butterflies was determined using the Checklist of Butterflies in the Philippine Islands (Treadaway 1995).

RESULTS AND DISCUSSION

Table 1. Status of butterflies of Maitum Village, Tandag, Surigao del Sur.

Species No.	Family/Species Name	Local	National	IUCN
A.	Family HESPERIIDAE			
1	<i>Aeromachus musca</i>	Very rare	Rare Philippine Endemic	Rare Endemic
2	<i>Ancistroides nigrita fumatus</i>	Very rare	Common	Common
3	<i>Borbo cinnara</i>	Very rare	Common	
4	<i>Halpe luteisquama</i>	Very rare	Common Philippine Endemic	
5	<i>H. sp.1</i>	Very rare	Undetermined	
6	<i>Isma feralia ferestrata</i>	Very rare	Rare Philippine Endemic	
7	<i>Notocrypta paralyso volux</i>	Rare	Common	Common Endemic
8	<i>Odontoptilum angulatunhelisa</i>	Very rare	Rare	
9	<i>Oriens californica</i>	Very rare	Rare Philippine Endemic	Common Endemic
10	<i>Parnara bada borneana</i>	Very rare	Rare	
11	<i>P. kawazoei</i>	Common	Common	
12	<i>Potanthus mingo mingo</i>	Very rare	Rare	Rare
13	<i>Pyronera liburnia minda</i>	Very rare	Rare Philippine Endemic	
14	<i>Tagiadesgana elegans</i>	Very rare	Common	Common
15	<i>T. japedustitus</i>	Rare	Common	Common
16	<i>T. trebellius martinus</i>	Very rare	Common	Common Philippine Endemic
17	<i>Taractrocera luzonensis</i>	Common	Common	
18	<i>Telicota sp.1</i>	Very rare	Undetermined	
B.	Family LYCAENIDAE			
19	<i>Acytolepis puspa bazilana</i>	Rare	Common	
20	<i>Allotinus fallaxa phacus</i>	Very rare	Common	
21	<i>Caleta angola angola</i>	Very rare	Rare	
22	<i>Catochrysops strabo luzonensis</i>	Very Common	Common	Philippine Endemic
23	<i>Celarchus archagathos archagathos</i>	Very rare	Rare Philippine Endemic	
24	<i>Cheritra orpheus orphmine</i>	Very rare	Common	
25	<i>Euchrysops cnejus cnejus</i>	Rare	Common	Common
26	<i>Everes lacturnus lacturnus</i>	Very Common	Common	
27	<i>Hypolycaena shirozui</i>	Very rare	Rare Philippine Endemic	

28	<i>H. sipylus tharrytas</i>	Rare	Common	
29	<i>Jamides celeno lydanus</i>	Very Common	Common	Rare
30	<i>J. cleodus cleodus</i>	Very rare	Common	
31	<i>J. philatu sosis</i>	Very rare	Rare	
32	<i>Lampides boeticus</i>	Very rare	Common	Common
33	<i>Nacaduba bereniceleei</i>	Rare	Rare	
34	<i>N. sp. 1</i>	Very rare	Undetermined	
35	<i>N. sp. 2</i>	Very rare	Undetermined	
36	<i>N. sp. 3</i>	Very rare	Undetermined	
37	<i>Pithecops corvus corax</i>	Very rare	Common	
38	<i>Prosotas dubiosa subardates</i>	Very rare	Rare, New record in Mindanao	
39	<i>P. sp. 1</i>	Very rare	Undetermined	
40	<i>Zizina otis oriens</i>	Very rare	Rare	
	Family NYMPHALIDAE			
41	<i>Acroptalmia albofasciata</i>	Very rare	Rare Mindanao Endemic	
42	<i>Amathusia phidippus pollicaris</i>	Rare	Common	Common
43	<i>A. sp.1</i>	Very rare	Undetermined	
44	<i>Anosia melanippus edmondii</i>	Very rare	Common	Common
45	<i>Athyma maenus semperi</i>	Very rare	Rare Rare Philippine	
46	<i>Discophora philipina</i>		Endemic	
47	<i>Dophla evelina proditrix</i>	Rare	Common Common Mindanao	
48	<i>Elymnias beza beza</i>	Very rare	Endemic	
49	<i>Euploe amulciber mindanensis</i>	Rare	Common	
50	<i>Euthalia alpheda cusama</i>	Very rare	Rare	
51	<i>Faunisphaon leucis</i>	Very rare	Common	
52	<i>Hypolimnas anomala anomala</i>	Rare	Common	Common
53	<i>H. bolina philippensis</i>	Very rare	Common	Common
54	<i>Junonia almata almata</i>	Very Common	Common	Common
55	<i>J. atlithes atlithes</i>	Rare	Common	
56	<i>J. hedonia ida</i>	Very Common	Common	Common
57	<i>Lasippae busa laetitia</i>	Very rare	Common	
58	<i>Lethe europa cevanna</i>	Very rare	Common	
59	<i>Lexias panopus miscus</i>	Very rare	Common	
60	<i>Melanitis atrax lucillus</i>	Very Common	Common Rare Philippine	
61	<i>M. boisduvalia boisduvalia</i>	Very rare	Endemic	
62	<i>M. leda leda</i>	Very Common	Common Rare Philippine	Common
63	<i>Moduza thespias</i>	Very rare	Endemic Rare Philippine	
64	<i>Mycalesis felderi felderi</i>	Rare	Endemic	
65	<i>M. janardana micromede</i>	Rare	Common	
66	<i>M. micromede micromede</i>	Rare	Rare	
67	<i>M. mineus philippina</i>	Common	Common	Rare
68	<i>M. sp.1</i>	Rare	Undetermined	
69	<i>M. tagala semirasa</i>	Common	Rare	
70	<i>Neptis cymela nitetus</i>	Very rare	Common	
71	<i>N. mindorana pseudosoma</i>	Rare	Common	
72	<i>N. pampanga boholic</i>	Very rare	Rare	Common
73	<i>Orsotriaena medus medus</i>	Rare	Common	
74	<i>Pantoporia dama commixta</i>	Very rare	Common	Common
75	<i>Rhinopalpa polynice validice</i>	Very rare	Common	
76	<i>Symbrenthia lilaea semperi</i>	Very rare	Common	Common

77	<i>Tanaecia leucotaenia quamarina</i>	Very rare	Common	
78	<i>Ypthima sempera chaboras</i>	Very Common	Common Philippine Endemic	Common Philippine Endemic
79	<i>Y. sempera sempera</i>	Rare	Common Philippine Endemic	
80	<i>Y. sensilis</i>	Very rare	Rare Philippine Endemic	
81	<i>Y. stellerastellera</i>	Very Common	Common Philippine Endemic	
82	<i>Zeuxidia amethystus amethystina</i>	Very rare	Rare	
D. Family PAPILIONIDAE				
83	<i>Arisbe decolor tigris</i>	Very rare	Very rare, Eastern Mindanao Endemic	
84	<i>A. euryplulus gordion</i>	Rare	Common	
85	<i>Atrophaneura semperi aphthonia</i>	Very rare	Rare	Rare Threatened, Philippine Endemic
86	<i>Grapium agamemnon agamemnon</i>	Very rare	Common	
87	<i>Lamproptera meges decius</i>	Very rare	Common	
88	<i>Menelaides deiphobus rumanzovia</i>	Very rare	Common	
89	<i>M. helenus hystaspes</i>	Very rare	Common Philippine Endemic	Common Philippine Endemic
90	<i>M. polytes ledebouria</i>	Very rare	Common	Common
91	<i>Pachliopta kotzebuea philippus</i>	Rare	Common	
92	<i>P. mariae mariae</i>	Very common	Common	
93	<i>Troides rhadamantus</i>	Very rare	Common Philippine Endemic	Common
E. Family PIERIDAE				
94	<i>Appia snepheleelis</i>	Rare	Rare	
95	<i>Catopsilia pyranthe pyranthe</i>	Common	Common	Common
96	<i>Cepora aspasia orantia</i>	Very rare	Common	
97	<i>Delias hyparete luzonensis</i>	Rare	Common	
98	<i>D. hyparete mindanensis</i>	Very rare	Common	
99	<i>Eurema alitha alitha</i>	Very common	Common Mindanao Endemic	Common Mindanao Endemic
100	<i>E. blanda valli volans</i>	Rare	Common	Common
101	<i>E. hecabeta miathis</i>	Very common	Common	
102	<i>E. saritalas arilata</i>	Very rare	Rare Philippine Endemic	Rare Philippine Endemic
103	<i>E. sp.1</i>	Very rare	Undetermined	
104	<i>Pareronia boebera trinobantes</i>	Very rare	Common	




Fig. 1. Map of Tandag City, Surigao del Sur (from: Municipal Development and Planning Office).




Fig. 2. Map of Maitum, Tandag, Surigao del Sur (Trinidad, 2004)



Fig. 3. Portion of agro-ecosystem, Maitum, Tandag, Surigao del Sur.



Fig. 4. Portion of dipterocarp forest, Maitum, Tandag, Surigao del Sur.



Fig. 6. Delineated 2 kilometer transect line in the agroecosystem of Maitum, Tandag, Surigao del Sur.



Fig. 6. Delineated 2 kilometer transect line in the dipterocarp forest of Maitum, Tandag, Surigao del Sur.



Fig. 7. Butterfly catching trap baited with overripe banana.



Fig. 8. Trap sampling using brightly colored clothes to attract butterflies to enhance species richness (adapted from Mohagan & Treadaway, 2010).




Fig. 9. Plot of abundance of butterflies of Maitum across two study stations.




Fig. 10. Shannon-Weiner plot of diversity for agroecosystem and dipterocarp forest sampling site




Fig. 11. Dendrogram on the cluster analysis of butterfly distribution in Maitum Village, Tandag, Surigao del Sur across the two sampling stations.



Fig. 12. 1a-8b. Species of Family Hesperidae



Fig. 13. 9a-16b. Species of Family HesperIIDae



Fig. 14. 17a-18b. Species of Family Hesperidae;
19a-24b. Species of Family Lycaenidae.



Fig. 15. 25a-32b. Species of Family Lycaenidae.



Fig. 16. 33a-40b. Species of Family Lycaenidae.



Fig. 17. 41a-48b. Species of Family Nymphalidae.



Fig. 18. 49a-56b. Species of Family Nymphalidae.



Fig. 19. 57a-64b. Species of Family Nymphalidae.



Fig. 20. 65a-72b. Species of Family Nymphalidae.



Fig. 21. 73a-80b. Species of Family Nymphalidae.



Fig. 22. 81a-82b. Species of Family Nymphalidae
83a-88b Species of Family Papilionidae



Fig. 23. 89a-93b. Species of Family Papilionidae
94-a-96-b. Species of Family Pieridae



Fig. 24. 97a-104b. Species of Family Pieridae

Species Composition

A total of 104 species of butterflies were collected in Maitum Village, Tandag, Suriagao del Sur. These species belong to 68 genera and 5 families of butterflies. Of these, eighteen (18) were from the family HesperIIDae, twenty two (22) were from the family Lycaenidae, forty (42) were from the family Nymphalidae, eleven (11) were from the family Papilionidae and eleven (11) were from the family Pieridae (Table 1). The result was lower as compared to the study of Mohagan and Treadaway (2010) in Mt. Hamiguitan, Davao Oriental with 142 species.

This may be due to the differences of the sampling techniques employed. The study of Mohagan and Treadaway (2010) employed an additional of two more methods of sampling, the Quadrat and Belt transect sampling method which were not employed in this study. There were species of butterflies common to both studies. This implies that butterflies have a certain degree of habitat preference. Butterflies select their habitat in relation to their food and host plants (Borkar & Komarpant 2004). Aside from, sampling techniques employed, the study site of Mohagan and Treadaway (2010) in Mt. Hamiguitan, Davao Oriental had a larger study site with greater number of habitats sampled and sampling duration was longer. Observation of Mohagan and Treadaway (2010) was also true to Cayabyab (2000) in Mt. Makiling, Laguna with 145 species. The study of Cayabyab (2000) used a modified version of Pollard's (1977) transect method that is similar to the method used in the current study.

The current study used 2 kilometer transects; although Cayabyab (2000) used 1 kilometer transects only, Cayabyab (2000) had a greater number of transects (7), than the current study (2). Another major difference between the current study, Cayabyab (2000) did not employ butterfly catching traps. The current study had a smaller sampling area than that of Cayabyab (2000). With reference to duration of study, Cayabyab (2000) had an exceedingly longer sampling duration. The greater sampling duration and the larger area in Mt. Makiling, Laguna could be a reason why the current study has a lesser number of species compared to Cayabyab (2000).

Although rapid surveys and extrapolation methods help provide a general framework for assessing biodiversity, understanding the

dynamics of species diversity in communities requires long term data sets that partition diversity into its natural components of space and time (De Vries & Walla 2001). It appears that methods, size of study site, and length of sampling period has a direct implication on the results of a study.

Compared to the results of Salmoy (2010) in Camiguin Island, the present study had a greater number of species than the former (81 species). In terms of methods, Salmoy (2010) had more sampling methods than the current study. In fact, methods used by Salmoy (2010) are identical to those of Mohagan and Treadaway (2010). With reference to sampling duration, the current study had a shorter duration, 2 months compared to the study of Salmoy (2010) with 6 months. With reference to study site, Salmoy (2010) in Camiguin Island had a larger area for sampling than the current study, although the sampling area of Salmoy (2010) is still considered a volcanic area. According to Weibull and Ostman (2003) landscape features largely affect species composition especially for the most mobile group of butterflies. A possible reason for the higher species composition of the current study would be the higher disturbance due to the volcanic activity in the area of Salmoy (2010) in Camiguin Island.

The current study had a greater number of species than Lit (2001) in Mount Banahao de Lucban, Quezon Province with 77 species. In terms of methods, the study of Lit (2001) and the current study were quite different due to the fact that the current study used more sampling methods. In terms of study site, the current study had smaller study site than Lit (2001). In terms of sampling duration, the study of Lit (2001) had a longer duration.

Aside from the difference in methods and duration of the study, butterfly composition differs from different study sites, which shows that geographical location influence butterfly composition. This observation is supported by Heaney (1991) and Mohagan and Treadaway(2010).Asper observation,vegetation typesalsoaffectspecies composition. The dipterocarp forest had higher species composition at 50% disconcondantwihagroecosystem. This observation is consistent to Salmoy (2010) and Mohagan and Treadaway (2010) where food plants availability differ from one vegetation type to another, so that butterfly composition also differs with vegetation types.

Species Diversity

Species Abundance

Adequacy of sampling was not reached for both sampling stations. This is mostly due to the limited sampling period and limited resources, not to mention the security condition in the study site. New species may be listed additionally if sampling will be extended.

In the current study, butterfly abundance was higher in dipterocarp forest (434) than in the agroecosystem (386) due to greater presence and diversity of food plants in the dipterocarp forest. This result is consistent with Torreno et al. (2007) in Mt Hamiguitan, Davao Oriental with a higher abundance in dipterocarp forest (288) than agroecosystem (89). According to Schneider (2003), both habitat characteristics and landscape structure influenced species numbers and abundance of butterflies in grasslands.

In the study of Mohagan and Treadaway (2010) in Mt Hamiguitan, Davao Oriental, the agroecosystem 1 (25) and agroecosystem 2 (26) had higher abundance than the dipterocarp 1 (15) and dipterocarp 2 (11). In Salmoy (2010), the agroecosystems of Mt. Timpoong (495) and Mt. Hibok-hibok (265) had higher abundance than the dipterocarp forests of both mountains (235). According to Mohagan and Treadaway (2010), abundance decreases with elevation.

It is interesting to note that in the studies of Mohagan and Treadaway (2010) and Salmoy (2010), there was a significant difference in the elevation of the agroecosystem and dipterocarp forest. In the current study, there was little difference in the elevation of the two study stations, therefore other factors like greater diversity of vegetation can be the reason for the greater abundance in the dipterocarp forest than agroecosystem and the type of habitat.

The agroecosystems studied by Mohagan and Treadaway (2010) and Salmoy (2010) were purely terrestrial but the agroecosystem of this present study was a ricefield with coconut trees. There were fewer types of plants in the agroecosystem of the current study than in the studies of Mohagan and Treadaway (2010) and Salmoy (2010). Melendez (2001) demonstrated that the existence of butterflies in one's place is influenced by reproductive season and presence of food plants.

Species Richness

The dipterocarp forest of Maitum had greater number of species (89) compared to the agroecosystem (51). The result of Mohagan and Treadaway (2010) for species richness is parallel to this current study in which there was an increasing trend in the agro ecosystems, dipterocarp and montane forests. Butterfly fauna may be affected and endangered by many factors including land use, forest cultivation and plant protection (Özden 2003). The anthropogenic use of land can be a factor to the number of species observed in the study. In other words, agricultural lands have a lesser species richness than the forested areas.

The species richness result of the current study is contradictory to Salmoy (2010) in which the agroecosystems of Mt. Timpoong (50) and Mt. Hibok- hibok (30) had higher species richness than dipterocarp forests (22). According to Schneider (2003), the amount of adjacent forest, flower abundance, field size and estimated nutrient levels are factors influencing butterfly species richness. Due to the security situation in the site of the current study, the landscape and flora of the dipterocarp forest were considerably kept intact. This may explain the greater species richness in the dipterocarp forest. Hansel et al. (2006) said that availability of host plants and food plants affects the presence of butterflies.

Shannon-Weiner Index for Diversity

In the current study, the average Shannon Weiner index of Maitum is low ($H' = 1.5$); when viewed in terms of study stations, the dipterocarp forest had fair ($H' = 1.638$) than the agroecosystem with a low level ($H' = 1.369$). This result is consistent with Ballentes et al. (2006) in Mt. Malindang, in which diversity was higher in forest ecosystems than in agroecosystems. Vegetation structure and taxonomic composition are well known to have a major influence on community diversity (MacArthur, Recher & Cody, 1966; Southwood et al., 1979). Extraction of trees, even at low densities can result to considerable damage to residual stand and severe declines in biodiversity (Johns, 1992; Kellman & Tackaberry, 1999 cited from Lewis, 2001). The reduction of tree density can be a factor for the lesser butterfly diversity of the agroecosystem compared to the dipterocarp forest.

The presence of bushes and trees had probably favored butterfly diversity by creating shelter and offering other resources (Schneider, 2003). It is interesting to note that the agroecosystem area had almost no naturally growing trees left; this can be the reason for the fair level of butterfly diversity of the dipterocarp forest.

The result of the current study is consistent with the results of Mohagan and Treadaway (2010) in which the agroecosystem had a slightly higher diversity value. The current study is contradictory to that of Salmoy (2010) in which the agroecosystem had high level of diversity. When compared to the study of Chiong (2007) in Gumata Falls, Magsal, Guinuyoran, Valencia City, the current study had a significantly lower diversity. This could be due to the sampling effort exerted by Chiong (2007) as it may be due to the varied plants available and disturbances in the area. In Salmoy (2010) the agroecosystem of Mt. Timpoong (1.70) and Mt. Hibok-hibok (1.48) has greater diversity than the dipterocarp of both mountains (1.34) in Camiguin Island. Her results are different from the current study.

Similarity Index

The cluster analysis of the species composition has shown that the agroecosystem and dipterocarp forest have low similarity of species composition with only 52.683%. This suggests that only 50% of the species are concordant to both the rest, and the remaining 50% are discordant. This result is parallel to the result of Salmoy (2010) in which there was clustering of the agroecosystem and dipterocarp forest in Mt. Timpoong, Camiguin de Mindanao. The similarity of the species composition of the agroecosystem and dipterocarp forest of the current study is due to the close proximity of the study stations in Maitum, Tandag, Surigaodel Sur. According to De Veries et al. (1997), apparent similarity among habitats is likely an area effect due to close proximity of the dipterocarp forest and recent human disturbance that created a patchwork of successional habitats in the agroecosystem.

Precipitation was present in both the summer and rainy season sampling periods, although the rainy season sampling period had more rainy days than the summer sampling period. Air temperature ranges from 28 - 40 °C, with no observable temperature differences in the two habitat types and during the two sampling periods. The lack

of difference of precipitation and air temperature could be a reason for the 50% concordant species observed in the study.

The difference in vegetation and in the two study stations can be a reason for the nearly 50% discordant butterfly species in the current study. This result is parallel to the results of Ballentes et al. (2006), in which little species similarities were observed between the vegetation types of Mt. Malindang.

Degree of disturbance in the two study stations can also be a reason for the nearly 50% discordant species of butterflies in Maitum Village, Tandag, Surigao del Sur, there might be forest specialist species present in Brangay Maitum, Tandag, Surigao del Sur. The greater disturbance in the agroecosystem can be a factor for the discordant species. According to Koh (2007), butterfly species restricted to undisturbed forests often have narrower geographical ranges than species found in disturbed habitats. Forest butterflies are more likely to be impacted by forest modification than those that are known to occur in disturbed habitats (Koh 2007).

Status

Danielsen and Treadaway (2004) reviewed the status of 915 species and 910 subspecies of butterflies known in the Philippines, of these there were 133 globally threatened and conservation-dependent Philippine endemic taxa. The Philippines is biodiversity giant, but is sad to note that the Philippines is also a biodiversity hotspot.

Of the 104 species sampled in the current study, 57 were common (54.8%), 16 were rare (15.3%), and 22 (21.1%) were endemics; 12 were rare Philippine endemics (11.5%), 6 were common Philippine endemics (5.7%), 2 common Mindanao endemics (1.9%), 1 rare Mindanao endemic (.96%), 1 very rare eastern Mindanao endemic (.96%), and 9 (8.6%) were undetermined.

The endemism percentage of the current study is lower than that of Mohagan & Treadaway (2010) (31%) in Mt. Hamiguitan. The study of Hansel et al. (2006) in Mt. Malindang, Misamis Occidental (62.5%) had higher percentage of endemism than the current study. The study of Salmoy (2010) in Camiguin Island, had similar endemism (22%) to the current study. In the study of Ballentes et al. (2006) in Mt. Malindang, there were two endemic butterflies, which was lower than the current study.

The rare Philippine endemics were: *Aeromachus musca*, *Isma feralia ferestrata*, *Oriens californica*, *Pyrroneura liburniaminda*, *Celarchusarchagathos sarchagathos*, *Hypolycaen ashirozui*, *Discophora philippina*, *Melanitis boisduvalia boisduvalia*, *Moduza thespias*, *Mycalesis felderi felderi*, *Ypthima sensilis*, and *Eurema sarilata sarilata*. Common Philippine endemics were: *Halpelutei squama*, *Ypthima sempera chaboras*, *Ypthima sempera sempera*, *Ypthima stelleria stelleria*, *Menelaides helenushystaspes*, and *Triodes rhadamantus*.

The common Mindanao endemics were: *Elymnas bezabeza* and *Eurema alithaalitha*. The rare Mindanao endemic species was *Acrophtalmia albofasciata* and the very rare Eastern Mindanao endemic species was *Arisbe decoloritigris*. There were 22 endemic species in Maitum Village, Tandag, Surigao del Sur, with 19 endemic species in the dipterocarp forest and 9 endemic species in the agroecosystem.

Noteworthy to mention among the species sampled in the current study is *Prosotas dubios asubardates*, a rare species which is also a new record in Mindanao. Other noteworthy samples are *Arisbedecoloritigris*, a very rare Eastern Mindanao endemic butterfly and *Acrophtalmia albofasciata*, a rare Mindanao endemic butterfly. Danielsen and Treadaway (2004) mentioned Sulu and Mindanao hold disproportionate numbers of threatened butterflies.

There is a great need for further biodiversity studies in Mindanao and Surigao del Sur since, numerous species may become extinct before they are even documented. During the conduct of the study, trunks of lumber floated on the river near the study site. According to Koh (2007), many Southeast Asian butterflies are endemic to the region and face the grim prospect of global extinction if current levels of deforestation were to continue. Butterflies are highly sensitive to habitat disturbance, and have been used as an indicator taxon for ecological research (Koh 2007). In addition to the small scale logging, sand and gravel is a common business in the current study area, this can be detrimental to the butterfly species present in the area.

Only a few species of butterflies had an IUCN conservation status, a few of these species are: *Mycalesis mineus philippina* (rare), *Atrophaneura semperi aphthonia* (rare, threatened, Philippine endemic), *Eurema alithaalitha* (common Mindanao endemic), *Eurema sarilata sarilata* (rare Philippine endemic). Data on the conservation status of

insects are scarce because the IUCN Red List is predisposed towards larger and better-studied taxa (Rodrigues et al. 2006). It is apparent that the IUCN system is difficult to use with regard to butterfly studies. A more comprehensive list for butterfly status is needed, although this requires an extensive compilation of numerous studies.

CONCLUSIONS AND RECOMMENDATIONS

Maitum, Tandag, Surigao del Sur is the home of one hundred four (104) species belonging to 5 families and 68 genera. Across the sampling stations, the dipterocarp forest showed a higher species abundance. The Shannon-Weiner index of diversity showed low level: Based on Kruger scale (2005), the dipterocarp forest of Maitum, Tandag, Surigao del Sur was at the fair level (1.638) of diversity than the agroecosystem area (1.369). The diversity level of Maitum is still low. Dendrogram index of similarity between the agroecosystem area and dipterocarp infers that the species of the two stations are 52.683% similar. This suggests that the two habitats sampled are both important for the conservation of butterflies in Maitum, Tandag, Surigao del Sur. The dipterocarp forest of Maitum, Tandag, Surigao del Sur is the home of 19 endemic species and the agroecosystem area is home of 9 endemic species for conservation.

The study recommends the use of other sampling techniques like belt transect and plot sampling and canopy nets to enhance species composition by sampling high flying or canopy butterflies. Further sampling should be done to augment species richness and in order to satisfy the species accumulation curve requirement and that other types of baits be used in the butterfly traps so as to improve species richness and status by sampling on the other areas of Tandag, Surigao del Sur.

ACKNOWLEDGMENTS

The authors are thankful to their families (Lorna, Bebiana, Lucio Ramirez; Susana, Roeland); (Dave, Dale Joy, Ard, John Dale and Dustin Kean Mohagan) for their financial and moral support. Grateful thanks are due to Dr. Lesley C. Lubos and Dr. Genaro V. Japos for their constant encouragements to disseminate our research results;

Central Mindanao University President for the support; Dr. Victor B. Amoroso and Dr. Florfe M. Acma for their valuable suggestions for the improvement of this paper.

LITERATURE CITED

- Ballentes, M.G., A.B. Mohagan, M.C.P. Espallardo, M.O. Zarcilla, and V.P. Gapud
2006 Arthropod faunal diversity and relevant interrelationships of critical resources in Mt. Malindang. Participatory Knowledge Generation: The Case of Biodiversity Research Programme (BRP) for Development, Mt. Malindang, Philippines. Southeast Asian Regional Center for Graduate Study and Research in Agriculture (SEARCA), Biodiversity Research Programme for Development in Mindanao: Focus on Mt. Malindang and Environs. pp 80, 86, 167.
- Borkar, M.R. and N. Komarpant
2004 Diversity, abundance, and habitat associations of butterfly species in Bondla Wildlife Sanctuary of Goa, India. *Zoos' Print Journal* , 19:10. pp 1648-1653.
- Cayabyab, B.F.
2000 A survey of Rhopalocera (Lepidoptera) of Mount Makiling, Laguna, Philippines. *The Philippine Entomologist*. 14:1. pp 105-106.
- Chiong, L.S.
2007 Diversity of butterflies in the Vicinity of Gumata Falls, Magsal, Guinuyoran, Valencia City, Bukidnon. (Thesis) CMU College of Arts and Sciences. College. Musuan, Bukidnon. pp 10 – 39.
- Danielsen, F. and C.G. Treadaway
2004 Priority conservation areas for butterflies (Lepidoptera: Rhopalocera) in the Philippine Islands. *Animal Conservation*. The Zoological Society of London. United Kingdom 10 (1017): 79-92.

De veries, P.J. and T.R. Walla

2001 Species diversity and community structure in neotropical fruit feeding butterflies. *Biological Journal of the Linnean Society* 42: (1 & 2) p. 1 Jan. 1991. [On line] Accessed on April 1, 2011. Available from <http://www.bio-nica.info/BibliotecaDeVries2001SpatialEffects.pdf>.

Hansel, C.G., A.G. Villarino, A.B. Mohagan, S.S. Nacua, T.O. Poblete, E.C. Roa, R.A. Seronay, F.M. Freire, B.A. Roscom, M.M. Gorospe, K.P. Culminas, W.C. Gay, D.G. Bacaltos

2006 Assessing the headwaters of Layawan River: Linkage between the terrestrial and aquatic ecosystem in Mt. Malindang, Misamis Occidental, Biodiversity Research Programme (BRP) for Development in Mindanao: Focus on Mt. Malindang and Environs. National Support Secretariat (NSS) SEAMEO SEARCA, College, Laguna 4031 Philippines.

Heaney, L.R.

1991 An analysis of patterns of distribution and species richness among Philippine fruit bats (Pteropodidae). *Bulletin of the American Museum of Natural History* 206: 145-167.

Kruger, M.A.

2005 The effects of insect diversity on three production fields: an apple orchard, a pepper field, and a garden. Sydney Australia. As cited by Mohagan A.B. 2007. *Diversity and Distribution of Butterflies in Mt. Hamiguitan Wildlife Sanctuary, Davao Oriental*, (Dissertation). CMU Graduate School. College. Musuan, Bukidnon.

Koh, L.P.

2007 Impacts of land use change on South- East Asian forest Butterflies: A review. *Journal of Applied Ecology* 44: 703-713.

Lewis, O.T.

2001 Effect of experimental selective logging on tropical butterflies. *Conservation Biology*. 15 (2): 389-400. [On line] Accessed on April 1, 2011. Available from <http://users.ox.ac>.

uk~zool0376Consbiolbelize.pdf.

Lit, I.L.

2001 A taxonomic list of butterflies (Lepidoptera: Papilionoidea and Hesperoidea) from Mount Banahao De Lucban, Quezon Province. *The Philippine Entomologist*. 15 (2): 151-152.

Macarthur, R.H., H. Recher, and M.L. Cody

1966 On the relation between habitat selection and species diversity. *American Naturalist* 100: 319-325.

Mendez, L.E.

2001 An inventory of butterflies in the vicinity of Lake Apo, Guinoyuran, Valencia Bukidnon. (Thesis).CMU College of Arts and Sciences.College.Musuan, Bukidnon. p 6.

Mohagan, A.B. and C.G. Treadaway

2010 Diversity and status of butterflies across vegetation types of Mt. Hamiguitan, Davao Oriental, Philippines. *Asian Journal of Biodiversity*. 1 (1): 1-24.

Municipal Planning AND Development Office (Tandag).

2004 Executive Summary. pp 1, 5, & 9.

Özden, Ö.

2003 Butterfly diversity and abundance in *Pinusbrutia* forests of Northern Cyprus. p 2. [On line] Accessed on April 1, 2011. Available from httpwww.undp-act.orgdataarticlesbdp_butterfly.pdf

POLLARD, E. and T.J. YATES

1995 Monitoring butterflies for ecology and conservation. Chapman and Hall St. Edmunds burry Press. Burry St. Edmunds, Suffolk, Great Britain.pp 14.

Rodrigues, A.S.L., J.D. Pilgrim., J.F. Lamoreux, Hoffman M. and T.M. Brooks.

2006 The value of the IUCN Red List for conservation. *Trends in*

Ecology and Evolution 21: 71-76 February 2006.

Salmoy, J.B.

2010 Diversity, status, and distribution of butterflies in Mt. Timpoong and Mt. Hibok-Hibok, Camiguin Island, Philippines (Graduate Thesis). CMU Graduate School. College. Musuan, Bukidnon pp 27-142.

Schneider, C.

2003 The influence of landscape structure on butterfly diversity and movement in grasslands- A comparison of two agricultural areas in Southern Sweden. (Dissertation). Swedish University of Agricultural Sciences, Alnarp. College pp 3- 40.

Southwood, T.R.E, V.K. Brown and P.M. Reader

1979 The relationship of plant and insect diversities in succession. Biological Journal of the Linnaean Society 12: 327-348 December 1979.

Torreno, H.S. Mohagan, A.B. Ramon, J.N. and H.C. Cahilog

2007 Survey and conservation of butterflies in Mt. Hamiguitan, Davao Oriental, p 4 Monograph.

Treadaway, C.G.

1995 Checklist of Butterfly in the Philippine Islands (Lepidoptera: Rhopalocera). Nachrichten des Entomologischen Vereins Apollo, Suppl. 14: 7-118; Frankfurt am Main Journal, Frankfurt, Germany.

Trinidad, E.T.

2004 Section Index Map: Maitum Village. Municipal Assessor's Office.

United Tourist Promotions.

2008 EZ Maps (tourist map) City of Tandag.

Warguez, D.A., K.M. Villamor, Tabaranza B.R. and M.J. Edrial
2009 Mammalian diversity in a logged over lowland forest of Mt.
Hilong- Hilong Range, Tandag, Surigao del Sur, Philippines.
WCSP 18th Annual Philippine Biodiversity Symposium
“Biodiversity in the Philippine Uplands” Symposium Manual
,April 15- 17, 2009. University of the Philippines, Baguio City.
pp 61.